

A Note On Cubic Splines, AMATH 352, March 4, 2002

We would like to use a spline to approximate a function represented by the points $(0, 0)$, $(1, 0)$, $(3, 2)$ and $(4, 2)$. The first task is to determine the spacing between the points h_k , the slopes d_k and then (though the solution of a system of equations) the second derivatives of the splines $s''(x_k) = m_k$.

$$\begin{aligned} h_0 &= x_1 - x_0 = 1 & h_1 &= x_2 - x_1 = 2 & h_2 &= x_3 - x_2 = 1 \\ d_0 &= \frac{y_1 - y_0}{x_1 - x_0} = 0 & d_1 &= \frac{y_2 - y_1}{x_2 - x_1} = 1 & d_2 &= \frac{y_3 - y_2}{x_3 - x_2} = 0 \end{aligned}$$

Next, set up the equations for the second derivatives (the m_k) which have been derived by requiring the continuity of the first and second derivatives of the splines at x_1 and x_2 :

$$\begin{aligned} 2(h_0 + h_1)m_1 + h_1m_2 &= 6(d_1 - d_0) - h_0m_0 \\ h_1m_1 + 2(h_1 + h_2)m_2 &= 6(d_2 - d_1) - h_2m_3 \end{aligned}$$

Note that there will be $N - 1$ of these equations when the spline interpolates $N + 1$ points. Since there are $N + 1$ m_k 's in this case, we need to apply the end conditions on the splines to determine m_0 and m_3 , the second derivatives at the end points. There are three options that we've introduced in this class:

- **Natural Spline** The second derivatives are zero at the ends:

$$\begin{aligned} m_0 &= 0 \\ h_0m_0 + 2(h_0 + h_1)m_1 + h_1m_2 &= 6(d_1 - d_0) \\ h_1m_1 + 2(h_1 + h_2)m_2 + h_2m_3 &= 6(d_2 - d_1) \\ m_3 &= 0 \end{aligned} \quad \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 6 & 2 & 0 \\ 0 & 2 & 6 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} m_0 \\ m_1 \\ m_2 \\ m_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 6 \\ -6 \\ 0 \end{pmatrix}$$

- **Clamped Spline** The first derivative of the spline is specified at the end points. Evaluating the first derivatives of the splines at x_0 and x_3 and requiring that they equal $f'(x_0)$ and $f'(x_3)$ yields the following system for the m_k :

$$\begin{aligned} 2h_0m_0 + h_0m_1 &= 6(d_0 - f'(x_0)) \\ h_0m_0 + 2(h_0 + h_1)m_1 + h_1m_2 &= 6(d_1 - d_0) \\ h_1m_1 + 2(h_1 + h_2)m_2 + h_2m_3 &= 6(d_2 - d_1) \\ h_2m_2 + 2h_2m_3 &= 6(f'(x_3) - d_2) \end{aligned} \quad \begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 6 & 2 & 0 \\ 0 & 2 & 6 & 1 \\ 0 & 0 & 1 & 2 \end{pmatrix} \begin{pmatrix} m_0 \\ m_1 \\ m_2 \\ m_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 6 \\ -6 \\ 0 \end{pmatrix}$$

- **Not-a-knot Spline** Without specifying any extra conditions at the end points (other than that the spline interpolates the data points there), the not-a-knot spline requires that the third derivative of the spline is continuous at x_1 and x_{N-1} . One can determine the equations for m_0 and m_N by requiring that $s'''_0(x_1) = s'''_1(x_1)$. (A similar condition holds at x_{N-1} .)

$$\begin{aligned} h_1m_0 - (h_0 + h_1)m_1 + h_0m_2 &= 0 \\ h_0m_0 + 2(h_0 + h_1)m_1 + h_1m_2 &= 6(d_1 - d_0) \\ h_1m_1 + 2(h_1 + h_2)m_2 + h_2m_3 &= 6(d_2 - d_1) \\ h_2m_1 - (h_1 + h_2)m_2 + h_1m_3 &= 0 \end{aligned} \quad \begin{pmatrix} 1 & -3 & 2 & 0 \\ 1 & 6 & 2 & 0 \\ 0 & 2 & 6 & 1 \\ 0 & 2 & -3 & 1 \end{pmatrix} \begin{pmatrix} m_0 \\ m_1 \\ m_2 \\ m_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 6 \\ -6 \\ 0 \end{pmatrix}$$

After solving the appropriate system for the end conditions that you have chosen, evaluate the coefficients of the splines:


$$s(x) = \begin{cases} s_0(x) = s_{0,0} + s_{0,1}(x-x_0) + s_{0,2}(x-x_0)^2 + s_{0,3}(x-x_0)^3 & x \in [x_0, x_1] \\ s_1(x) = s_{1,0} + s_{1,1}(x-x_1) + s_{1,2}(x-x_1)^2 + s_{1,3}(x-x_1)^3 & x \in [x_1, x_2] \\ s_2(x) = s_{2,0} + s_{2,1}(x-x_2) + s_{2,2}(x-x_2)^2 + s_{2,3}(x-x_2)^3 & x \in [x_2, x_3] \end{cases}$$

where $s_{k,0} = y_k$ $s_{k,1} = d_k - \frac{h_k}{6}(2m_k + m_{k+1})$ $s_{k,2} = \frac{m_k}{2}$ $s_{k,3} = \frac{m_{k+1} - m_k}{6h_k}$.

With natural end conditions, I found that $m_0 = m_3 = 0$, $m_1 = \frac{3}{2}$ and $m_2 = -\frac{3}{2}$, so that:

$$s(x) = \begin{cases} s_0(x) = 0 - \frac{1}{4}(x-0) + 0(x-0)^2 + \frac{1}{4}(x-0)^3 & x \in [0, 1] \\ s_1(x) = 0 + \frac{1}{2}(x-1) + \frac{3}{4}(x-1)^2 - \frac{1}{4}(x-1)^3 & x \in [1, 3] \\ s_2(x) = 2 + \frac{1}{2}(x-3) - \frac{3}{4}(x-3)^2 + \frac{1}{4}(x-3)^3 & x \in [3, 4] \end{cases}$$

Here is a plot of the natural spline that passes through these four points. At right is a closeup of the splines between $x = 3$ and $x = 4$ to highlight the difference in the end conditions. Note that the spline appears not to have any curvature at the end points, while the not-a-knot spline does have a non-zero second derivative at the right end in this case. The clamped spline is constructed to have zero slope at the end points.



Note that the spline function in MATLAB computes a not-a-knot spline by default. If $X = [0 \ 1 \ 3 \ 4]$ and $Y = [0 \ 0 \ 2 \ 2]$, the not-a-knot spline can be computed and plotted in MATLAB with `plot(x,ppval(spline(X,Y),x))`. Specifying additional data points at the beginning and end of the interval will give a clamped spline with those extra values as the slopes at the endpoints of the intervals. The command `plot(x,ppval(spline(X,[0 Y 0]),x))` would give the clamped spline plotted here with $f'(0) = f'(4) = 0$.